

TAGITOL™ V (BARIUM SULFATE) ORAL SUSPENSION, 40% w/v

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TAGITOL V safely and effectively. See full prescribing information for TAGITOL V.

TAGITOL V (barium sulfate) oral suspension

Initial U.S. Approval: 2016

INDICATIONS AND USAGE

TAGITOL V is a radiographic contrast agent indicated in adult patients for use in computed tomography (CT) colonography as a fecal tagging agent (1)

DOSAGE AND ADMINISTRATION

- The recommended dose is:
 - One 20 mL bottle (8g barium sulfate) with each meal (breakfast, lunch and dinner) the day before the CT colonography examination (2.1).
 - Total dose = 3 bottles (24 g barium sulfate)
- For oral use only (2.2).

DOSAGE FORMS AND STRENGTHS

- Oral suspension: barium sulfate (40% w/v) 20 mL single dose bottles as a ready to use suspension for oral administration (3)

CONTRAINDICATIONS

TAGITOL V is contraindicated in patients with:

- Known or suspected perforation of the gastrointestinal (GI) tract (4)
- Known obstruction of the GI tract (4)

- Conditions associated with high risk of GI perforation or aspiration (4)
- Known hypersensitivity to barium sulfate or any of the excipients of TAGITOL V (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity reactions: Emergency equipment and trained personnel should be immediately available (5.1)
- Intra-abdominal barium leakage: May occur in conditions which increase the risk of perforation such as - carcinoma, GI fistula, inflammatory bowel disease, gastric or duodenal ulcer, appendicitis, diverticulitis, or severe stenosis or obstructing lesions of the GI tract (5.2)
- Delayed GI transit and obstruction: Patients should maintain adequate hydration in days following a barium sulfate procedure to avoid obstruction or impaction by baroliths (5.3)
- Aspiration pneumonitis: Caution is recommended in patients with a history of food aspiration and in patients with known swallowing disorders (5.4)

ADVERSE REACTIONS

Common adverse reactions include nausea, vomiting, diarrhea and abdominal cramping (6)

To report SUSPECTED ADVERSE REACTIONS, contact Bracco Diagnostics Inc at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

See 17 for PATIENT COUNSELING INFORMATION

Revised: 8/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

<p>1 INDICATIONS AND USAGE</p> <p>2 DOSAGE AND ADMINISTRATION</p> <p>2.1 Recommended Dosage</p> <p>2.2 Important Administration Instructions</p> <p>3 DOSAGE FORMS AND STRENGTHS</p> <p>4 CONTRAINDICATIONS</p> <p>5 WARNINGS AND PRECAUTIONS</p> <p>5.1 Hypersensitivity Reactions</p> <p>5.2 Intra-abdominal Barium Leakage</p> <p>5.3 Delayed Gastrointestinal Transit and Obstruction</p> <p>5.4 Aspiration Pneumonitis</p> <p>5.5 Systemic Embolization</p> <p>6 ADVERSE REACTIONS</p> <p>8 USE IN SPECIFIC POPULATIONS</p> <p>8.1 Pregnancy</p> <p>8.2 Lactation</p>	<p>8.4 Pediatric Use</p> <p>8.5 Geriatric Use</p> <p>11 DESCRIPTION</p> <p>12 CLINICAL PHARMACOLOGY</p> <p>12.1 Mechanism of Action</p> <p>12.2 Pharmacodynamics</p> <p>12.3 Pharmacokinetics</p> <p>13 NONCLINICAL TOXICOLOGY</p> <p>13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility</p> <p>16 HOW SUPPLIED/STORAGE AND HANDLING</p> <p>16.1 How Supplied</p> <p>16.2 Storage and Handling</p> <p>17 PATIENT COUNSELING INFORMATION</p>
---	---

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

TAGITOL V is indicated for use in adult patients for use in computed tomography (CT) colonography as a fecal tagging agent.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

- The recommended oral dose of TAGITOL V is one 20 mL bottle (8 g barium sulfate) with each meal (breakfast, lunch and dinner) the day before the colonography examination. Total dose = 3 bottles (24 g barium sulfate).

2.2 Important Administration Instructions

- TAGITOL V is typically provided to the patient for self-administration. Advise patients to carefully read and follow the Patient Instructions for Use to be provided to the patient.
- Shake bottle for 15 seconds prior to administration.
- For oral use only.
- Encourage patients to hydrate following the barium sulfate procedure.
- Discard any unused suspension.

3 DOSAGE FORMS AND STRENGTHS

Oral suspension: barium sulfate (40% w/v) supplied as a ready-to-use suspension in a 20 mL, single-dose, plastic bottle for oral administration. Each 20 mL bottle contains 8 g of barium sulfate.

4 CONTRAINDICATIONS

TAGITOL V is contraindicated in patients with:

- known or suspected perforation of the gastrointestinal (GI) tract;
- known obstruction of the GI tract;
- high risk of GI perforation such as those with a recent GI perforation, acute GI hemorrhage or ischemia, toxic megacolon, severe ileus, post GI surgery or biopsy, acute GI injury or burn, or recent radiotherapy to the pelvis;
- high risk of aspiration such as those with prior aspiration, tracheo-esophageal fistula, or obtundation;
- known hypersensitivity to barium sulfate or any of the excipients of TAGITOL V.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Barium sulfate preparations contain a number of excipients, including natural and artificial flavors and may induce serious hypersensitivity reactions. The manifestations include hypotension, bronchospasm and other respiratory impairments, dermal reactions including rashes, urticaria and itching. A history of bronchial asthma, atopy, or a previous reaction to a contrast agent may increase the risk for hypersensitivity reactions. Emergency equipment and trained personnel should be immediately available for treatment of a hypersensitivity reaction.

5.2 Intra-abdominal Barium Leakage

The use of TAGITOL V is contraindicated in patients at high risk of perforation of the GI tract [see *Contraindications (4)*]. Administration of TAGITOL V may result in leakage of barium from the GI tract in the presence of conditions that increase the risk of perforation such as carcinomas, GI fistula, inflammatory bowel disease, gastric or duodenal ulcer, appendicitis, or diverticulitis, and in patients with a severe stenosis at any level of the gastrointestinal tract, especially if it is distal to the stomach. The barium leakage has been associated with peritonitis and granuloma formation.

5.3 Delayed Gastrointestinal Transit and Obstruction

Orally administered barium sulfate may accumulate proximal to a constricting lesion of the colon, causing obstruction or impaction with development of baroliths (inspissated barium associated with feces) and may lead to abdominal pain, appendicitis, bowel obstruction, or rarely perforation. Patients with the following conditions are at higher risk for developing obstruction or baroliths: severe stenosis at any level of the GI tract, impaired gastrointestinal motility, electrolyte imbalance, dehydration, on a low residue diet, on medications that delay GI motility, constipation, cystic fibrosis, Hirschsprung disease, and the elderly. To reduce the risk of delayed GI transit and obstruction, patients should maintain adequate hydration following a barium sulfate procedure.

5.4 Aspiration Pneumonitis

The use of TAGITOL V is contraindicated in patients at high risk of aspiration [see *Contraindications (4)*]. Oral administration of barium is associated with aspiration pneumonitis, especially in patients with a history of food aspiration or with compromised swallowing mechanism. Vomiting following oral administration of barium sulfate may lead to aspiration pneumonitis.

5.5 Systemic Embolization

Barium sulfate products may occasionally intravasate into the venous drainage of the large bowel and enter the circulation as a "barium embolus" leading to potentially fatal complications which include systemic and pulmonary embolism, disseminated intravascular coagulation, septicemia and prolonged severe hypotension. Although this complication is exceedingly uncommon after oral administration of barium sulfate products, monitor patients for potential intravasation when administering barium sulfate.

6 ADVERSE REACTIONS

The following adverse reactions have been identified from spontaneous reporting or clinical studies of barium sulfate administered orally. Because the reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to drug exposure:

- Nausea, vomiting, diarrhea and abdominal cramping.
- Serious adverse reactions and fatalities include aspiration pneumonitis, barium sulfate impaction, intestinal perforation with consequent peritonitis and granuloma formation, vasovagal and syncopal episodes.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

TAGITOL V is not absorbed systemically following oral administration, and maternal use is not expected to result in fetal exposure to the drug [see *Clinical Pharmacology (12.3)*].

8.2 Lactation

Risk Summary

TAGITOL V is not absorbed systemically by the mother following oral administration, and breastfeeding is not expected to result in fetal exposure to the drug [see *Clinical Pharmacology (12.3)*].

8.4 Pediatric Use

TAGITOL V is not indicated for pediatric use.


8.5 Geriatric Use

Clinical studies of TAGITOL V do not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

11 DESCRIPTION

TAGITOL V (barium sulfate) is a radiographic contrast agent that is supplied as a 40% w/v, off-white to lightly colored, free-flowing, ready-to-use suspension with an apple aroma for oral administration. The active ingredient barium sulfate is designated chemically as

BaSO₄ with a molecular weight of 233.4 g/mol and the following chemical structure:



TAGITOL V contains the following excipients: carboxymethylcellulose sodium, citric acid, glycerin, maltodextrin, natural and artificial apple flavor, polysorbate 80, potassium sorbate, purified water, saccharin sodium, simethicone emulsion, sodium benzoate, sodium citrate, xanthan gum, and xylitol.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Due to its high atomic number, barium (the active ingredient in TAGITOL V) is opaque to x-rays and therefore acts as a positive contrast agent for radiographic studies.

12.2 Pharmacodynamics

Barium sulfate is biologically inert and has no known pharmacological effects.

12.3 Pharmacokinetics

Under physiological conditions, barium sulfate passes through the gastrointestinal tract in an unchanged form and is absorbed only in pharmacologically insignificant amounts.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No animal studies have been performed to evaluate the carcinogenic potential of barium sulfate or potential effects on fertility.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

TAGITOL V (barium sulfate) is an oral suspension (40% w/v) supplied as a box of three 20 mL HDPE bottles. Each bottle contains 8 grams barium sulfate.

Provided as: 24 boxes, each containing 3 (20 mL) bottles (NDC 32909-814)

16.2 Storage and Handling

Store at USP controlled room temperature 20 to 25°C (68 to 77° F). Protect from freezing.

17 PATIENT COUNSELING INFORMATION

After administration, advise patients to:

- Maintain adequate hydration [see *Dosage and Administration (2.2)* and *Warnings and Precautions (5.3)*].
- Seek medical attention for worsening of constipation or slow gastrointestinal passage [see *Warnings and Precautions (5.3)*].
- Seek medical attention for any delayed onset of hypersensitivity: rash, urticaria, or respiratory difficulty [see *Warnings and Precautions (5.1)*].

Rx only

Manufactured by
EZEM Canada Inc
Anjou (Quebec) Canada H1J 2Z4

For
Bracco Diagnostics Inc.
Monroe Township, NJ 08831