

ProHance[®] Multipack[™] (Gadoteridol) Injection, 279.3 mg/mL

Pharmacy Bulk Package - Not for Direct Infusion

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PROHANCE MULTIPACK safely and effectively. See full prescribing information for PROHANCE MULTIPACK.

PROHANCE MULTIPACK (gadoteridol injection), for intravenous use PHARMACY BULK PACKAGE -- NOT FOR DIRECT INFUSION
Initial U.S. Approval: 2003

WARNING: NEPHROGENIC SYSTEMIC FIBROSIS
See full prescribing information for complete boxed warning

Gadolinium-based contrast agents (GBCAs) increase the risk for NSF among patients with impaired elimination of the drugs.

Avoid use of GBCAs in these patients unless the diagnostic information is essential and not available with non-contrasted MRI or other modalities. NSF may result in fatal or debilitating systemic fibrosis affecting the skin, muscle and internal organs.

The risk for NSF appears highest among patients with:

- chronic, severe kidney disease (GFR less than 30 mL/min/1.73m²), or
- acute kidney injury

Screen patients for acute kidney injury and other conditions that may reduce renal function. For patients at risk for chronically reduced renal function (e.g. age greater than 60 years, hypertension or diabetes), estimate the glomerular filtration rate (GFR) through laboratory testing (5.1).

RECENT MAJOR CHANGES

Indications and Usage, MRI of the Central Nervous System (1.1) 12/2020

INDICATIONS AND USAGE

ProHance Multipack is a gadolinium-based contrast agent indicated for magnetic resonance imaging (MRI) to visualize:

- lesions with disrupted blood brain barrier and/or abnormal vascularity in the brain (intracranial lesions), spine and associated tissues in adults and pediatric patients, including term neonates (1.1)
- lesions in the head and neck in adults (1.2)

DOSAGE AND ADMINISTRATION

- Dispense multiple single doses into separate sterile syringes for intravenous administration (2.3)
- Recommended dose in adult and pediatric patients is 0.2 mL/kg (0.1 mmol/kg) body weight administered as rapid intravenous infusion or bolus (2.1)
- Follow injection with a saline flush of at least 5 mL normal saline (2.1)

DOSAGE FORMS AND STRENGTHS

Injection: contains 279.3 mg/mL (0.5 mmol/mL) of gadoteridol supplied in a pharmacy bulk pack (2.3, 3, 16)

CONTRAINDICATIONS

Allergic or hypersensitivity reactions to ProHance (4)

WARNINGS AND PRECAUTIONS

- Nephrogenic Systemic Fibrosis has occurred in patients with impaired elimination of GBCAs. Higher than recommended dosing or repeated dosing appears to increase risk (5.1)
- Hypersensitivity: anaphylactic/anaphylactoid reactions with cardiovascular, respiratory and cutaneous manifestations, ranging from mild to severe reactions including shock can occur. Monitor patients closely for need of emergency cardiorespiratory support (5.2)
- Gadolinium is retained for months or years in brain, bone, and other organs. (5.3)

ADVERSE REACTIONS

The most commonly reported adverse reactions are nausea and taste perversion with an incidence \geq 0.9% (6.1)

To report SUSPECTED ADVERSE REACTIONS, Contact Bracco Diagnostics Inc. at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

USE IN SPECIFIC POPULATIONS

Pregnancy: Use only if imaging is essential during pregnancy and cannot be delayed. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 12/2020

FULL PRESCRIBING INFORMATION: CONTENTS* WARNING: NEPHROGENIC SYSTEMIC FIBROSIS

- 1 INDICATIONS AND USAGE
 - 1.1 MRI of the Central Nervous System (CNS)
 - 1.2 MRI of Extracranial/Extraspinal Head and Neck
- 2 DOSAGE AND ADMINISTRATION
 - 2.1 Recommended Dose
 - 2.2 Administration
 - 2.3 Directions for Proper Use of Pharmacy Bulk Package
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Nephrogenic Systemic Fibrosis (NSF)
 - 5.2 Hypersensitivity Reactions
 - 5.3 Gadolinium Retention
 - 5.4 Acute Kidney Injury (AKI)
- 6 ADVERSE REACTIONS
 - 6.1 Clinical Trials Experience
 - 6.2 Post-marketing Experience

8. USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Renal Impairment
 - 10 OVERDOSAGE
 - 11 DESCRIPTION
 - 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
 - 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
 - 14 CLINICAL STUDIES
 - 14.1 MRI of the CNS
 - 14.2 MRI of the Head and Neck
 - 16 HOW SUPPLIED/STORAGE AND HANDLING
 - 17 PATIENT COUNSELING INFORMATION
- *Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNING: NEPHROGENIC SYSTEMIC FIBROSIS
Gadolinium-based contrast agents (GBCAs) increase the risk for NSF among patients with impaired elimination of the drugs. Avoid use of GBCAs in these patients unless the diagnostic information is essential and not available with non-contrasted MRI or other modalities. NSF may result in fatal or debilitating systemic fibrosis affecting the skin, muscle and internal organs.

The risk for NSF appears highest among patients with:

- chronic, severe kidney disease (GFR less than 30 mL/min/1.73m²), or
- acute kidney injury

Screen patients for acute kidney injury and other conditions that may reduce renal function. For patients at risk for chronically reduced renal function (e.g. age greater than 60 years, hypertension or diabetes), estimate the glomerular filtration rate (GFR) through laboratory testing.

For patients at highest risk for NSF, do not exceed the recommended ProHance dose and allow a sufficient period of time for elimination of the drug from the body prior to re-administration [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

1.1 MRI of the Central Nervous System (CNS)

ProHance is indicated for magnetic resonance imaging (MRI) in adults and pediatric patients including term neonates to visualize lesions with disrupted blood brain barrier and/or abnormal vascularity in the brain (intracranial lesions), spine and associated tissues.

1.2 MRI of Extracranial/Extraspinal Head and Neck

ProHance is indicated for MRI in adults to visualize lesions in the head and neck.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose

The recommended dose for adult and pediatric patients, including term neonates, is 0.2 mL/kg (0.1 mmol/kg) administered as a rapid intravenous infusion (10 mL/min to 60 mL/min) or bolus (greater than 60 mL/min). Table 1 provides weight-adjusted recommended dose volumes.

Body Weight (kg)	Volume to be Administered (mL)
2.5	0.5
5	1
10	2
20	4
30	6
40	8
50	10
60	12
70	14
80	16
90	18
100	20
110	22
120	24
130	26
140	28
150	30

MRI of the CNS in Adults

- A supplementary dose of 0.4 mL/kg (0.2 mmol/kg) may be given up to 30 minutes after the first dose in adult patients with normal renal function suspected of having poorly visualized CNS lesions, in the presence of negative or equivocal scans
- The safety and efficacy of supplementary dosing have not been established in pediatric patients

2.2 Administration

- Visually inspect ProHance for particulate matter and discoloration prior to use
- Do not administer the solution if it is discolored or particulate matter is present
- Concurrent medications or parenteral nutrition should not be physically mixed with contrast agents and should not be administered in the same intravenous line because of the potential for chemical incompatibility
- Inject at least a 5 mL normal saline flush immediately after ProHance injection to ensure complete administration
- Imaging procedures should be completed within 1 hour

2.3 Directions for Proper Use of Pharmacy Bulk Package NOT FOR DIRECT INFUSION

The pharmacy bulk package is used as a multiple dose container with an appropriate transfer device to fill empty sterile syringes. Use the following procedures when transferring ProHance from the pharmacy bulk package to individual syringes:

- Use of this product is restricted to a suitable work area, such as a laminar flow hood, utilizing aseptic technique
- Prior to entering the vial, remove the seal and cleanse the rubber closure with a suitable antiseptic agent
- The container closure may be penetrated only one time, utilizing a suitable transfer device or dispensing set that allows measured dispensing of the contents
- Once the pharmacy bulk package is punctured, it should not be removed from the aseptic work area during the entire period of use
- Withdrawal of container contents should be accomplished without delay. A maximum time of 8 hours from initial closure entry is permitted to complete fluid transfer operations
- Any unused contents must be discarded by 8 hours after initial puncture of the bulk package
- Once drawn into syringe, administer transferred agent promptly for single-dose administration

3 DOSAGE FORMS AND STRENGTHS

ProHance Multipack is supplied as a sterile, nonpyrogenic, and colorless to slightly yellow solution available in 50 mL pharmacy bulk packages for intravenous administration. Each mL contains 279.3 mg (0.5 mmol/mL) of gadoteridol for injection.

4 CONTRAINDICATIONS

ProHance is contraindicated in patients with known allergic or hypersensitivity reactions to ProHance [see Warnings and Precautions (5.2)].

5 WARNINGS AND PRECAUTIONS

5.1 Nephrogenic Systemic Fibrosis (NSF)

Gadolinium-based contrast agents (GBCAs) increase the risk for NSF among patients with impaired elimination of the drugs. Avoid use of GBCAs among these patients unless the diagnostic information is essential and not available with non-contrast MRI or other modalities. The GBCA-associated NSF risk appears highest for patients with chronic, severe kidney disease (GFR less than 30 mL/min/1.73m²) as well as patients with acute kidney injury. The risk appears lower for patients with chronic, moderate kidney disease (GFR 30-59 mL/min/1.73m²) and little, if any, for patients with chronic, mild kidney disease (GFR 60-89 mL/min/1.73m²). NSF may result in fatal or debilitating fibrosis affecting the skin, muscle and internal organs. Report any diagnosis of NSF following ProHance Multipack administration to Bracco Diagnostics (1-800-257- 5181) or FDA (1-800-FDA-1088 or www.fda.gov/medwatch).

ProHance Multipack is indicated for acute kidney injury and other conditions that may reduce renal function. Features of acute kidney injury consist of rapid (over hours to days) and usually reversible decrease in kidney function, commonly in the setting of surgery, severe infection, injury or drug-induced kidney toxicity. Serum creatinine levels and estimated GFR may not reliably assess renal function in the setting of acute kidney injury. For patients at risk for chronically reduced renal function (for example, age greater than 60 years, diabetes mellitus or chronic hypertension), estimate the GFR through laboratory testing. Among the factors that may increase the risk for NSF are repeated or higher than recommended doses of a GBCA and the degree of renal

impairment at the time of exposure. Record the specific GBCA and the dose administered to a patient. For patients at highest risk for NSF, do not exceed the recommended ProHance dose and allow a sufficient period of time for elimination of the drug prior to re-administration. For patients receiving hemodialysis, physicians may consider the prompt initiation of hemodialysis following the administration of a GBCA in order to enhance the contrast agent's elimination. The usefulness of hemodialysis in the prevention of NSF is unknown. [see *Clinical Pharmacology (12)*].

5.2 Hypersensitivity Reactions

Anaphylactic and anaphylactoid reactions have been reported, involving cardiovascular, respiratory, and/or cutaneous manifestations. Some patients experienced circulatory collapse and died. In most cases, initial symptoms occurred within minutes of ProHance administration and resolved with prompt emergency treatment. Prior to ProHance administration, ensure the availability of trained personnel and medications to treat hypersensitivity reactions. Consider the risk for hypersensitivity reactions, especially in patients with a history of hypersensitivity reactions or a history of asthma or other allergic disorders. If such a reaction occurs, stop ProHance and immediately begin appropriate therapy. Observe patients for signs and symptoms of a hypersensitivity reaction during and for up to 2 hours after ProHance administration.

5.3 Gadolinium Retention

Gadolinium is retained for months or years in several organs. The highest concentrations (nanomoles per gram of tissue) have been identified in the bone, followed by other organs (e.g. brain, skin, kidney, liver, and spleen). The duration of retention also varies by tissue and is longest in bone. Linear GBCAs cause more retention than macrocyclic GBCAs. At equivalent doses, retention varies among the linear agents with Omniscan (gadodiamide) and Optimar (gadoversetamide) causing greater retention than other linear agents [Eovist (gadoxetate disodium), Magnevist (gadopentetate dimeglumine), MultiHance (gadobenate dimeglumine)]. Retention is lowest and similar among the macrocyclic GBCAs [Dotarem (gadoterate meglumine), Gadavist (gadobutrol), ProHance (gadoteridol)].

Consequences of gadolinium retention in the brain have not been established. Pathologic and clinical consequences of GBCA administration and retention in skin and other organs have been established in patients with impaired renal function [see Warnings and Precautions (5.1)]. There are rare reports of pathologic skin changes in patients with normal renal function. Adverse events involving multiple organ systems have been reported in patients with normal renal function without an established causal link to gadolinium retention [see Adverse Reactions (6.2)].

While clinical consequences of gadolinium retention have not been established in patients with normal renal function, certain patients might be at higher risk. These include patients requiring multiple lifetime doses, pregnant and pediatric patients, and patients with inflammatory conditions. Consider the retention characteristics of the agent when choosing a GBCA for these patients. Minimize repetitive GBCA imaging studies, particularly closely spaced studies when possible.

5.4 Acute Kidney Injury (AKI)

In patients with chronically reduced renal function, acute kidney injury requiring dialysis has occurred with the use of GBCAs. The risk of acute kidney injury may increase with increasing dose of the contrast agent; administer the lowest dose necessary for adequate imaging.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the prescribing information:

- Nephrogenic systemic fibrosis [see Boxed Warning and Warnings and Precautions (5.1)].
- Hypersensitivity reactions [see Contraindications (4) and Warnings and Precautions (5.2)].

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The adverse events described in this section were observed in clinical trials involving 3174 subjects (including 2896 adults and 278 pediatric subjects ages 0 to 17 years) exposed to ProHance. Approximately 48% of the subjects were men and ethnic distribution was 78% Caucasian, 6% Black, 3% Hispanic, 6% Asian, and 2% other. In 5% of the subjects, race was not reported. Average age was 47 years (range from 1 day to 91 years) and the exposure ranged from 0.03 to 0.3 mmol/kg. Overall, approximately 5.8% of subjects reported one or more adverse reactions during a follow-up period that ranged from 24 hours to 7 days after ProHance administration.

Table 2 lists adverse reactions that occurred in \geq 0.4% subjects who received ProHance.

Reaction	Rate (%) N = 3174
Nausea	1.4%
Dysgeusia	0.9%
Headache	0.7%
Dizziness	0.4%
Urticaria	0.4%

The following additional adverse events occurred in fewer than 0.4% of the subjects:

General disorders and administration site conditions:	Asthenia; chest discomfort, facial edema, feeling hot, injection site coldness, injection site erythema, injection site pain, injection site warmth, pain, pyrexia
Cardiac:	Angina pectoris, palpitations, atrio-ventricular block first degree
Ear and labyrinth disorders:	Ear discomfort, tinnitus
Eye disorders:	Eyed pruritus, injection increased
Gastrointestinal disorders:	Abdominal discomfort, abdominal pain, diarrhea, dry mouth, gingival pain, oral pruritis, swollen tongue, vomiting
Infections and infestations:	Gingivitis, rhinitis
Investigations:	Alanine aminotransferase increased, aspartate aminotransferase increased, blood chloride increased, blood pressure immeasurable, blood urea increased, hemoglobin decreased, heart rate decreased
Metabolism and nutrition disorders:	Decreased appetite, hypoglycemia
Musculoskeletal and connective tissue disorders:	Back pain, musculoskeletal stiffness
Nervous system disorders:	Formication, hypoesthesia, hypokinesia, lethargy, loss of consciousness, migraine, parosmia, presyncope, seizure, syncope, taste disorder
Psychiatric disorders:	Anxiety, mental status changes
Respiratory, thoracic and mediastinal disorders:	Cough, dry throat, dyspnea, nasal discomfort, throat irritation
Skin and subcutaneous tissue disorders:	Hyperhidrosis, pruritis, rash, rash morbilliform
Vascular disorders:	Flushing, hypotension, peripheral coldness, vascular rupture, vasodilatation, vasospasm

6.2 Post-marketing Experience

The following adverse reactions have been identified during post approval use of ProHance that were not observed in the clinical trials. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

MEDICATION GUIDE
PROHANCE® (prō-'han(t)s)
(Gadoteridol injection)
for intravenous use

What is PROHANCE?

- PROHANCE is a prescription medicine called a gadolinium-based contrast agent (GBCA). PROHANCE, like other GBCAs, is used with a magnetic resonance imaging (MRI) scanner.
- An MRI exam with a GBCA, including PROHANCE, helps your doctor to see problems better than an MRI exam without a GBCA.
- Your doctor has reviewed your medical records and has determined that you would benefit from using a GBCA with your MRI exam.

What is the most important information I should know about PROHANCE?

- PROHANCE contains a metal called gadolinium. Small amounts of gadolinium can stay in your body including the brain, bones, skin and other parts of your body for a long time (several months to years).
- It is not known how gadolinium may affect you, but so far, studies have not found harmful effects in patients with normal kidneys.
- Rarely, patients have reported pains, tiredness, and skin, muscle or bone ailments for a long time, but these symptoms have not been directly linked to gadolinium.
- There are different GBCAs that can be used for your MRI exam. The amount of gadolinium that stays in the body is different for different gadolinium medicines. Gadolinium stays in the body more after Omniscan or Optimark than after Eovist, Magnevist, or MultiHance. Gadolinium stays in the body the least after Dotarem, Gadavist, or ProHance.
- People who get many doses of gadolinium medicines, women who are pregnant and young children may be at increased risk from gadolinium staying in the body.
- Some people with kidney problems who get gadolinium medicines can develop a condition with severe thickening of the skin, muscles and other organs in the body (nephrogenic systemic fibrosis). Your healthcare provider should screen you to see how well your kidneys are working before you receive PROHANCE.

Do not receive PROHANCE if you have had a severe allergic reaction to PROHANCE.

Before receiving PROHANCE, tell your healthcare provider about all your medical conditions, including if you:

- have had any MRI procedures in the past where you received a GBCA. Your healthcare provider may ask you for more information including the dates of these MRI procedures.
- are pregnant or plan to become pregnant. It is not known if PROHANCE can harm your unborn baby. Talk to your healthcare provider about the possible risks to an unborn baby if a GBCA such as PROHANCE is received during pregnancy
- have kidney problems, diabetes, or high blood pressure
- have had an allergic reaction to dyes (contrast agents) including GBCAs

What are the possible side effects of PROHANCE?

- See “What is the most important information I should know about PROHANCE?”
- **Allergic reactions. PROHANCE can cause allergic reactions that can sometimes be serious. Your healthcare provider will monitor you closely for symptoms of an allergic reaction.**

The most common side effects of PROHANCE include: nausea, distortion of the sense of taste, and headache.

These are not all the possible side effects of PROHANCE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of PROHANCE.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. You can ask your healthcare provider for information about PROHANCE that is written for health professionals.

What are the ingredients in PROHANCE?

Active ingredient: gadoteridol

Inactive ingredients: calteridol calcium, tromethamine

Manufactured by: BIPSO GmbH-78224 Singen (Germany)

Manufactured for: Bracco Diagnostics Inc., Monroe Township, NJ 08831

For more information, go to www.imaging.bracco.com or call 1-800-257-5181.